

## JPE 2018 table of errata

As of January 31, 2022

### Monographs

| Page | Line  | Correction  | Error   |
|------|-------|---|---|
| 243  | ↑ 3   | 0.70 mL of 0.01 mol/L hydrochloric acid VS  | 0.07 mL of 0.01 mol/L hydrochloric acid VS  |
| 250* | ↑ 5   | <b>Absorbance</b>   | <b>Optical rotation</b>   |
| 468* | ↓ 12  | for ferric <u>salt</u> and for chloride   | for ferric <u>chloride</u> and for chloride   |
| 589  | ↓ 2-3 | <b>(2) Arsenic</b> – Prepare the test solution with 2.0 g of Methacrylic Acid Copolymer LD according to <u>Method 3</u> , and perform the test (not more than 1 ppm). | <b>(2) Arsenic</b> – Prepare the test solution with 2.0 g of Methacrylic Acid Copolymer LD according to <u>Method 2</u> , and perform the test (not more than 1 ppm). |
| 785* | ↑ 9   | and shake for about 15 <u>seconds</u> ,   | and shake for about 15 <u>minutes</u> ,   |
| 786  | ↑ 15  | <b>Viscosity</b> 350 – 550 <u>mm<sup>2</sup>/s</u> (Method 1, 20°C).  | <b>Viscosity</b> 350 – 550 <u>mm<sup>2</sup>s</u> (Method 1, 20°C).   |
| 836* | ↓ 10  | To <u>20 g</u> of Pregelatinized Starch   | To <u>2.0 g</u> of Pregelatinized Starch  |
| 858  | ↓ 12  | of which value is not <u>less</u> than 1.8% and not <u>more</u> than 2.2%.  | of which value is not <u>more</u> than 1.8% and not <u>less</u> than 2.2%.  |

\*: correction for the first print run.